

Question:

$$\text{Let: } x + y + z = 1$$

$$x^2 + y^2 + z^2 = 4$$

$$x^3 + y^3 + z^3 = 9$$

$$\text{Solve for } x^4 + y^4 + z^4$$

For extra credit, solve for the general case where:

$$a = x + y + z$$

$$b = x^2 + y^2 + z^2$$

$$c = x^3 + y^3 + z^3$$

Solution:

Let's jump right in and start with the general "extra credit" case.

$$\text{Let } d = x^4 + y^4 + z^4$$

$$\text{We can get to } d \text{ with } a^4 = (x+y+z)^4 = x^4 + y^4 + z^4 + 4*(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + 6*(x^2y^2 + x^2z^2 + y^2z^2) + 12*(x^2yz + xy^2z + xyz^2)$$

You can see there are lots of mixed terms in that expression we will have to evaluate.

Let's start at the bottom and work our way up. Evaluating  $a^2$  is a good place to start.

$$a^2 = (x + y + z)^2 =$$

$$x^2 + y^2 + z^2 + 2*(xy + xz + yz) =$$

$$b + 2(xy + xz + yz) \text{ -- Recall: } b = x^2 + y^2 + z^2$$

$$(1) xy + xz + yz = (a^2 - b)/2$$

$$ab = (x + y + z)*(x^2 + y^2 + z^2)$$

$$= x^3 + y^3 + z^3 + xy^2 + xz^2 + yx^2 + yz^2 + zx^2 + zy^2$$

$$= c + xy^2 + xz^2 + yx^2 + yz^2 + zx^2 + zy^2 \text{ -- Recall: } c = x^3 + y^3 + z^3$$

$$(2) xy^2 + xz^2 + yx^2 + yz^2 + zx^2 + zy^2 = ab - c$$

$$a^3 = (x + y + z)^3 =$$

$$x^3 + y^3 + z^3 + 3(xy^2 + xz^2 + yx^2 + yz^2 + zx^2 + zy^2) + 6xyz =$$

$$c + 3(ab - c) + 6xyz$$

$$6xyz = a^3 - c - 3(ab - c)$$

$$6xyz = a^3 + 2c - 3*ab$$

$$(3) xyz = (a^3 + 2c - 3ab)/6$$

$$b^2 = (x^2 + y^2 + z^2)^2 = x^4 + y^4 + z^4 + 2*(x^2y^2 + x^2z^2 + y^2z^2)$$

$$(4) x^2y^2 + x^2z^2 + y^2z^2 = (b^2 - d)/2 \quad \text{-- Recall: } d = x^4 + y^4 + z^4$$

$$\begin{aligned} a^2 * b &= (x^2 + y^2 + z^2 + 2*(xy + xz + yz)) * (x^2 + y^2 + z^2) = \\ &x^4 + y^4 + z^4 + 2*(x^2y^2 + x^2z^2 + y^2z^2) + 2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + 2(x^2yz + xy^2z \\ &+ xyz^2) = \\ &d + 2*(x^2y^2 + x^2z^2 + y^2z^2) + 2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + 2xyz*(x+y+z) = \\ &d + 2*(x^2y^2 + x^2z^2 + y^2z^2) + 2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + a(a^3 + 2c - 3ab)/3 = \\ &d + (b^2 - d) + 2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + a(a^3 - 3ab + 2c)/3 = \\ &b^2 + 2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + a(a^3 - 3ab + 2c)/3 \end{aligned}$$

$$2(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) = a^2 * b - b^2 - a(a^3 - 3ab + 2c)/3$$

$$x^3y + x^3z + y^3x + y^3z + z^3x + z^3y = a^2 * b/2 - b^2/2 - a^4/6 + a^2b/2 - ac/3$$

$$(5) x^3y + x^3z + y^3x + y^3z + z^3x + z^3y = a^2 * b - b^2/2 - a^4/6 - ac/3$$

$$\begin{aligned} a^4 &= (x + y + z)^4 = \\ &x^4 + y^4 + z^4 + 4*(x^3y + x^3z + y^3x + y^3z + z^3x + z^3y) + 6*(x^2y^2 + x^2z^2 + y^2z^2) + 12*(x^2yz + \\ &xy^2z + xyz^2) = \\ &d + 4*(a^2b - b^2/2 - a^4/6 - ac/3) + 6((b^2 - d)/2) + 12xyz(x+y+z) = \\ &d + 4a^2b - 2b^2 - (2/3)a^4 - (4/3)ac + 3b^2 - 3d + 12((a^3 + 2c - 3ab)/6)*a = \\ &d + 4a^2b - 2b^2 - (2/3)a^4 - (4/3)ac + 3b^2 - 3d + 2a^4 + 4ac - 6a^2b = \\ &4/3a^4 + (8/3)ac - 2a^2b + b^2 \end{aligned}$$

$$2d = 1/3a^4 + (8/3)ac - 2a^2b$$

$$d = a^4/6 + (4/3)ac - a^2b + b^2/2$$

Recall in the original problem,  $a = 1$ ,  $b = 4$ , and  $c = 9$ :

$$d = 1^4/6 + (4/3)1^2*9 - 1^2*4 + 4^2/2 = 97/6 = 16.166666.$$

Further reading:

Newton identities: [https://en.wikipedia.org/wiki/Newton's\\_identities](https://en.wikipedia.org/wiki/Newton's_identities)